

Literatur

zum Beitrag „Symptome managen in der Onkologie“ von Matthias Hellberg-Naegele

PflegeZeitschrift (74) 7/2021, S. 26-29

Armstrong, T. S. (2003). Symptoms Experience: A Concept Analysis. *Oncology Nursing Forum*, 30(4), 601–606. <https://doi.org/10.1188/03.ONF.601-606>

Armstrong, T. S. (2014). 2013 Special Topics Conference: Peaks and pitfalls in longitudinal studies of symptom outcome data. *Nursing Outlook*, 62(5), 305–312. <https://doi.org/10.1016/j.outlook.2014.05.005>

Dodd, M., Janson, S., Facione, N., Faucett, J., Froelicher, E. S., Humphreys, J., Lee, K., Miaskowski, C., Puntillo, K., & Rankin, S. (2001). Advancing the science of symptom management. *Journal of advanced nursing*, 33(5), 668–676.

Katz, S., Ford, A. B., Moskowitz, R. W., Jackson, B. A., & Jaffe, M. W. (1963). Studies of illness in the aged: The index of ADL: a standardized measure of biological and psychosocial function. *Jama*, 185(12), 914–919.

Kirsch, M., Halter, J., Stüssi, G., Dobbels, F., & De Geest, S. (2011). Patient Reported Outcomes in view of symptom experience of late effects and self-management of adult long-term survivors after allogeneic haematopoietic stem cell transplantation – A mixed methods study. PROVIVO.

Lawton, M. P., & Brody, E. M. (1969). Assessment of older people: Self-maintaining and instrumental activities of daily living. *The gerontologist*, 9(3_Part_1), 179–186.

Leppla, L., Naegele, M., Mößner, U., Rebafka, A., & Hasemann, M. (2016). Advanced practice nurses' activities in oncology and future perspectives—Improving patients' symptom burden as leading goal. Abteilungsworkshop, Saig.

Leppla, L., Naegele, M., Rebafka, A., & Koller, A. (2016). Orale Kryotherapie bei Hochdosis-Melphalan vor autologer Stammzelltransplantation. *Onkologische Pflege*, 6(1), 40–46.

McDaniel, R., & Rhodes, V. A. (1995). Symptom experience. *Seminars in Oncology Nursing*, 4(11), 232–234. [https://doi.org/10.1016/s0749-2081\(05\)80002-6](https://doi.org/10.1016/s0749-2081(05)80002-6)

Naegele, M., Leppla, L., Rebafka, A., Hasemann, M., & Engelhardt, M. (2017). Die Entwicklung von Symptommanagement-Strategien als Tätigkeitsfeld von APN am Beispiel von Diarrhoe bei Patienten mit Multiplem Myelom unter Hochdosis-Chemotherapie Melphalan. *Advanced Practice Nurses Magazin*, 1, 75–76.

Naegele, M., Kirsch, M., Ihorst, G., Fierz, K., Engelhardt, M., & De Geest, S. (2018). Symptom experience of multiple myeloma (syMMex) patients treated with autologous stem cell transplantation following high-dose melphalan: A descriptive longitudinal study. *Supportive Care in Cancer*, 26(3), 833–841. <https://doi.org/10.1007/s00520-017-3897-z>

National Cancer Institute. (2017). Common Terminology Criteria for Adverse Events (CTCAE) Version 5.0.

https://ctep.cancer.gov/protocoldevelopment/electronic_applications/docs/CTCAE_v5_Quick_Reference_5x7.pdf

National Cancer Institute. (2020). NCI-PRO-CTCAE Items German.
https://healthcaredelivery.cancer.gov/pro-ctcae/pro-ctcae_german.pdf

Rhodes, V. A., & Watson, P. M. (1987). Symptom distress—The concept: Past and present. *Seminars in Oncology Nursing*. 3(4), 242–247.

Simms, S. G., Rhodes, V. A., & Madsen, R. W. (1993). Comparison of prochlorperazine and lorazepam antiemetic regimens in the control of postchemotherapy symptoms. *Nursing research*, 42(4), 234–239.